IJCRT.ORG ISSN : 2320-2882
INTERNATIONAL JOURNAL OF CREATIVE RESEARCH THOUGHTS (IJCRT)
An International Open Access, Peer-reviewed, Refereed Journal

Biochemistry, Nutritional Composition, And Health Benefits Of Spinach (*Spinacia Oleracea*): A Comprehensive Review

S. Moorthi[1]
[1]Professor, Sree Sastha Arts and Science College, Chembarambakkam, Chennai, Tamil Nadu, India

Abstract: The leafy green spinach (Spinacia oleracea) is high in nutrients and has a solid reputation for promoting general health. Rich in vital vitamins, minerals, antioxidants, and other bioactive substances, spinach can help prevent and treat a number of chronic illnesses, such as cancer, heart disease, and metabolic problems. This paper takes a close look at the biochemistry, nutritional makeup, and health benefits of spinach. It also explores how different factors such as food processing methods and bioavailability to impact its nutritional value. Special attention is given to the plant's metabolic pathways, enzymatic activities, and secondary metabolites that contribute to its therapeutic potential. Finally, the paper highlights the need for future research aimed at improving nutrient absorption and maximizing the health benefits of spinach through better agricultural and processing techniques.

Keywords: Nutritional, Bioavailability, Spinach, Chronic Illness, Enzymatic activities

1. INTRODUCTION

Spinach (Spinaciaoleracea), a species belonging to the Amaranthaceae family, is extensively cultivated as a leafy green vegetable renowned for its remarkable nutritional profile and associated health benefits. Originating in Western and Central Asia, spinach has achieved worldwide acclaim owing to its versatile applications in both culinary arts and medicinal practices. Commonly consumed in diverse forms ranging from raw salads to cooked dishes, spinach is esteemed for providing an abundant array of essential nutrients, antioxidants, and phytochemicals.

With its abundance of minerals, vitamins, antioxidants, and other potent compounds, water spinach has recently attracted a lot of attention. [1]. Studies show that the stem, in particular, is loaded with crude protein, healthy fats, and fiber, making it a nutritious option that's available all year round [1]. Adding more water spinach into pelleted animal feed has even been shown to boost how well the animals digest organic matter, hinting at its value as a fiber-rich ingredient [2].

The leaves are just as impressive. They're easy to mix into a wide variety of dishes and are popular across all age groups. Even better, they bring a healthy dose of calcium, fiber, and iron to the table, enhancing the nutritional value of meals [3]. In fact, more and more creative ways are being explored to blend water spinach into traditional foods, showing just how versatile it can be [3].

Beyond just water spinach, spinach varieties in general are renowned for having elevated amounts of protein, iron, riboflavin, and vitamin C — making them an essential food source, especially for communities that might not have access to a wide range of nutritious options [4].

When it comes to farming, water spinach is pretty tough. It thrives in slightly acidic to neutral soils (pH 5 to 7) [5], loves warm temperatures above 23.9°C, and grows best in clay or marshy soil that's rich in organic matter and stays moist [5]. There's even evidence that its nutrient content goes up when it's grown in nutrient-rich water [6]. Farmers have a few different ways of growing it: some scatter seeds on raised beds and harvest monthly, others plant in neat rows for quicker, weekly harvests, and some grow it floating directly on water [3], [5], [7].

A leafy green vegetable with high nutritional value is spinach, has been gaining interest for its potential use in aquaponics systems where it can help with nutrient cycling and water quality improvement [8]. In aquaponics, growing water spinach alongside fish farming creates a synergistic system where plants absorb nutrients from fish waste, helping to clean the water and reduce organic material [9].

Researchers have studied the growth and productivity of water spinach in aquaponic circulatory systems, focusing on how well it absorbs mineral nutrients from fish waste and leftover feed [9], [10]. Water spinach is also being explored as a cost-effective substitute for fish meal in aquaculture because it offers an affordable, easy-to-cultivate source of plant protein [5].

Different parts of water spinach offer unique nutritional benefits. The stem contains higher concentrations of crude protein and fiber, while the leaf holds significant amounts of carbohydrates [1]. Although hydroponic methods show promise for cultivating water spinach, challenges remain in optimizing its growth and productivity across different stages of development [10].

To monitor the progress of water spinach, growers use data-driven techniques such as weekly measurements of plant height, leaf number, and fresh weight, along with qualitative assessments of overall plant development [11]. When grown with koi fish in aquaponic systems, water spinach has demonstrated notable productivity, showing increased stem length and leaf production. It has also contributed to better fish survival rates and absolute growth [9].

Aquaponics, which combines aquaculture and hydroponics, uses wastewater from fish farming to nourish plants. This process reduces the need for irrigation and supplies vital nutrients for plant growth [12]. The integrated system enhances sustainability by converting fish waste into valuable nutrients for plants, leading to better water conservation and efficient nutrient recycling [9], [13].

Getting the balance of nutrients between fish and plants right is essential to maintain a healthy system where nutrient generation and removal are in harmony [14]. Aquaponics also naturally improves water quality because the plant roots and beneficial bacteria around them help remove harmful substances like ammonia, which can be toxic to fish [15], [16].

Compared to traditional and hydroponic farming methods, aquaponics offers more advantages by reusing wastewater, enhancing production and profits by optimizing energy, water, and nutrient utilization, and supporting plant and ecosystem health [17]. The integration of plants, fish, and bacteria into a cohesive system is crucial for keeping every component healthy and ensuring high water quality [18].

In aquaponics systems, incorporating nutrient cycles offers a sustainable way to minimize waste and reduce the environmental impacts of food production, including excessive water use and nutrient losses [19]. In these systems, water circulates between the plant bed and the fish habitat, conserving water much more efficiently than traditional irrigation methods [20]. Adding plants into the aquaponic setup helps remove nutrients from the water, especially ammonia, which is toxic to fish. This not only improves water quality but also reduces the need for water disposal [16].

This integrated approach brings together aquaculture and hydroponics, using less water while supporting the sustainable co-cultivation of plants and fish [21], [22]. Over time, the hydroponic component of an aquaponic system also becomes a thriving home for microbiota that assist with biofiltration [22], [23], [24]. These beneficial microorganisms enhance water purification by removing harmful chemicals like ammonia and nitrites, boosting system health and stability.

To achieve best results from this integrated farming method, producers need to carefully plan aspects such as system design, fish stocking densities, plant densities, and fish feeding rates.

This comprehensive review aims to offer a detailed analysis of the biochemistry, nutritional composition, and wide-ranging health benefits of spinach by integrating the latest research findings. As more people turn toward plant-based diets, growing importance in understanding spinach's broad range of bioactive chemicals, along with the mechanisms through which they support human health.

The purpose of this study is to clarify, up-to-date overview of the nutritional composition, biochemical processes, and health benefits associated with spinach (*Spinaciaoleracea*). It aims to bring together recent research findings to help deepen our understanding of how spinach's bioactive compounds support health. The paper also explores spinach's role in sustainable farming systems like aquaponics and outlines the key factors that influence how well our bodies can absorb and use its nutrients. A further objective is to highlight future directions for research to enhance spinach's nutritional value and therapeutic use.

2. Spinach Botany and Agriculture

Spinach, formally known as Spinaciaoleracea, is an Amaranthaceae leafy green vegetable recognized for its nutritional value and cooking versatility [25]. Originating from ancient Persia, spinach has a long history of cultivation, with evidence suggesting it was grown as early as the 6th century AD before spreading to other parts of the world [26]. The plant usually grows up to about 30 centimeters tall and produces broad, succulent leaves that can be either smooth or crinkled depending on the variety.

Today, spinach is cultivated extensively around the world, with major growing regions including China, the United States, and Europe. It thrives best in temperate climates with well-drained soils. Several different varieties of spinach are grown, each offering unique characteristics such as differences in leaf texture, growth rate, and disease resistance. Incorporating spinach into aquaponics systems—an integrated approach to food production that blends hydroponics (growing plants without soil) and aquaculture (raising fish)—has gained popularity recently.

Spinach is widely appreciated for its culinary flexibility and impressive nutritional content. In recent years, it has gained more recognition as a functional food. It is a fast-growing, cool-season vegetable that prefers temperatures between 10°C and 20°C. Known for its broad, green, succulent leaves, spinach grows best in fertile, well-drained soils. It is generally classified into three main types: smooth-leafed, savoy (curled), and semi-savoy varieties. Smooth-leafed types are often used in processed foods, while savoy varieties are preferred for fresh consumption due to their textured leaves [26].

The nutrient composition of spinach is significantly influenced by cultivation methods. Temperature, light exposure, and soil quality are examples of environmental variables that might affect the amount of vitamins and minerals in the leaves. Using proper cultivation techniques can help maximize the bioavailability of essential nutrients in spinach, offering significant benefits to both consumers and the agricultural industry.

3. Biochemistry of Spinach

3.1 Photosynthesis and Metabolic Pathways

With the aid of chlorophyll pigments, photosynthesis transforms carbon dioxide and water into glucose and oxygen, hence influencing the nutritional makeup of spinach. Environmental variables including temperature, carbon dioxide concentration, and light intensity affect how effective this process is. Together, these factors determine the overall biomass and nutritional quality of spinach [27].

Chlorophyll and carotenoids are two important pigments found in spinach. Advanced methods including UV-visible spectroscopy, thin layer chromatography, and column chromatography can be used to identify and examine them [28]. These methods allow for the precise quantification and characterization of the pigments, which are essential for both photosynthesis and antioxidant defense.

Spinach participates in a number of important metabolic processes in addition to photosynthesis, including as glycolysis, the Krebs cycle, and the electron transport chain. These pathways contribute to energy production and the synthesis of important metabolites that support the plant's growth and nutritional value. Figure 1 shows Spinach Energy Metabolism.



Figure 1 Spinach Energy Metabolism

3.2 Bioactive Compounds

The numerous health advantages of spinach are attributed to its abundance of bioactive substances, which include vitamins, minerals, antioxidants, and other phytonutrients. It offers vital minerals including iron, calcium, and magnesium as well as vitamins A, C, and K, all of which are necessary for a number of bodily physiological processes.

Flavonoids, phenolic acids, and carotenoids are also found in spinach; these compounds have potent antioxidant qualities that aid in shielding cells from oxidative damage brought on by free radicals [29]. Depending on the growing conditions of various spinach species, these phytochemical profiles can change. [26].

Advanced analytical methods are needed to identify and quantify these bioactive substances. For accurate analysis, techniques including spectrophotometry, gas chromatography-mass spectrometry, and high-performance liquid chromatography are frequently employed.

3.3 Amino Acids

Vegetables are important sources of amino acids, which are vital for assessing their nutritional quality. Recent research has demonstrated a direct correlation between the light environment in which plants develop and the metabolism of amino acids [30].

A number of procedures are often involved in the analysis of amino acids in spinach, such as extraction, derivatization, and separation using methods such as gas chromatography-mass spectrometry or liquid chromatography-mass spectrometry.

Spinach displays complex metabolic pathways that regulate the production of many essential compounds. These include the electron transport chain, which produces a proton gradient needed for ATP synthesis; the Krebs cycle, which oxidizes acetyl-CoA to produce ATP; and glycolysis, which breaks down glucose to produce energy. The leaves, where photosynthesis takes place, are rich in chlorophyll and carotenoids that are essential for absorbing light.

Understanding these metabolic processes is critical for improving spinach cultivation and enhancing its nutritional value [31].

3.4 Primary and Secondary Metabolites

Plants produce both primary and secondary metabolites, which are essential for their growth, development, and interactions with the environment [32]. Primary metabolites, such as carbohydrates, lipids, and proteins, are directly involved in vital processes, while secondary metabolites contribute to defense, signaling, and adaptation. Secondary metabolites found in plants have various uses, such as their utilization in traditional medicines and pharmaceuticals [33]. Phenolic phytochemicals, a complex group of secondary metabolites, are responsible for adding flavor and color to fruits, vegetables, and grains [34].

Plants synthesize a variety of organic compounds known as secondary metabolites, which are crucial for protection against herbivores and environmental challenges [35]. These compounds hold significant pharmacological importance, and their structures have been described in over 100,000 instances [36]. The production of these metabolites is significantly influenced by growth and environmental factors [37], [38].

Alkaloids, flavonoids, phenolic compounds, and molecules containing nitrogen and sulfur are examples of secondary metabolites that naturally defend against infections, herbivores, and oxidative stress [39]. They are typically biosynthesized in specific cell types during certain life stages and generally accumulate in smaller amounts compared to primary metabolites [40]. Over time, plants have evolved to produce a diverse array of small molecular weight secondary metabolites that enhance survival and reproduction [41].

Terpenoids, derived from terpenes, are among the volatile constituents known for their protective qualities. These specialized metabolic pathways create molecules with functions that improve adaptability and survival. Unlike primary metabolites, which are necessary for basic life functions, secondary metabolites contribute to enhancing a plant's fitness and defense, often produced in limited quantities during specific developmental stages [42], [43].

Secondary metabolites produced by microorganisms, such as phenolics, terpenes, and substances containing nitrogen, also play important roles in shielding plants from biotic and abiotic stresses [44]. These specialized metabolites exhibit a remarkable diversity in structure, function, and biosynthetic pathways, with more than 2,140,000 secondary metabolites identified, divided into five main categories: alkaloids, nonribosomal polypeptides, fatty acid-derived compounds and polyketides, terpenoids and steroids, and enzyme cofactors [45].

Advancements in molecular techniques have made it possible to alter how secondary metabolites are expressed in plants [46]. Many of these compounds possess bioactive properties that can protect against a variety of diseases by interacting with specific cellular targets and modulating biochemical pathways [47]. They are also recognized for their toxicity and repellent properties, deterring herbivores and microbial threats [46].

The biological functions of plants, such as defense and interaction with other species, are supported by the structural diversity of natural products made from them [48]. Plants also produce volatile compounds, a subset of secondary metabolites, to defend themselves directly by repelling herbivores or indirectly by attracting predators of their attackers [49].

Secondary metabolites not only play a vital role in plant defense but also serve as valuable natural resources for medicine and agriculture [50], [51]. Their presence highlights the ability of plants to adapt to their environment and ensure survival [43], [52]. Emissions of volatile organic molecules, mostly terpenoids, from these interactions serve as crucial signals between plants and other living things, including pollinators, seed dispersers, herbivores, predators, parasitoids, and nearby plants [53].

Plant extracts, which are complex mixtures of general leaf volatiles and specific chemical components, have demonstrated repellency against pests [54]. Since plants cannot physically escape herbivores, they have evolved to produce low molecular weight chemicals that are either toxic or repellent to animals [55]. These chemical defenses can reduce palatability, act as poisons, or enhance the expression of defense-related genes [56].

Secondary metabolites contribute significantly to plant defense by mediating the discharge of environmental protecting substances [56], [57]. Many of these substances also function as antimicrobial agents, anticancer agents, analgesics, antidiarrheal agents, and antifungal substances, making them highly valuable in medical applications [58].

Gaining molecular insights into the proteins and genes engaged in secondary metabolite production can aid in metabolic engineering to strengthen plant tolerance to herbivores [59]. Plants react to herbivore attacks through morphological, biochemical, and molecular changes [60]. Additionally, defense compounds from ancestral plant varieties can be restored through breeding or transferred across different species [61].

3.3 Enzymatic Activity

Enzymes are fundamental to various metabolic processes in plants, including photosynthesis, respiration, and nutrient assimilation. They regulate the biosynthesis of essential metabolites, promoting plant growth and enhancing responses to environmental stimuli. Enzymes also contribute to plant resistance against insects by producing repellent or toxic compounds, inhibiting insect digestive enzymes, or strengthening plant cell walls [60], [62]. These strategies can function independently or work together synergistically [60]. Understanding the synthesis, regulation, and breakdown of plant specialized metabolites is critical for improving their use in medicines, insecticides, and other valuable products.

Plant enzymes are central to major metabolic activities such as photosynthesis, respiration, and nutrient assimilation [63]. Supporting plant growth, development, and adaptation to environmental conditions depends on these mechanisms [64]. Numerous chemicals that aid plants in fending off herbivores and environmental stress are produced through secondary metabolism, which is regulated by enzyme activity [65]. The kinds and quantities of metabolites generated are mostly determined by enzymes, which improve plant growth and responses to environmental stimuli.

Insect resistance in plants is closely linked to enzyme activities that lead to the production of defensive compounds, the inhibition of insect digestive processes, or the reinforcement of plant structural barriers [66]. These defense strategies may act independently or in combination to protect the plant.

The regulation of both primary and secondary metabolite biosynthesis by enzymes is crucial for optimizing plant development and defense mechanisms [67]. Enzymes drive secondary metabolic pathways, catalyzing the formation of a diverse array of compounds that serve to protect the plant from herbivores and environmental stressors [68]. However, more research is needed to identify additional factors involved in defense responses and to fully characterize the underlying molecular mechanisms [69].

In addition to their role in metabolic processes like photosynthesis and respiration, enzymes are indispensable for nutrient assimilation. Specialized metabolites produced under enzymatic control offer significant defensive advantages against herbivores and environmental pressures [70]. The influence of enzymes on metabolite production is key to improving plant resilience and growth under varying environmental conditions.

Enzymes involved in glucose and amino acid metabolism, redox balance, stress responses, photosynthesis, signal transduction, and protein processing are often studied through proteomics to gain a better understanding of their roles and molecular mechanisms [71]. A deeper understanding of how plant specialized metabolites are synthesized, regulated, and catabolized is vital for enhancing their applications in medicine, insecticide development, and other high-value industries [67], [72], [73], [74].

3.5 Bioavailability and Absorption

Bioavailability, or the degree and speed at which these substances enter the bloodstream and reach their intended site of action, is a key factor in the therapeutic efficacy of plant metabolites. The chemical structure of the metabolite, its stability and solubility, the herbal product's composition, and the physiological traits of the person ingesting it are some of the variables that affect bioavailability [75]. The ability of plant metabolites to exert their beneficial effects is shaped by these complex interactions.

Metabolic pathways within the gastrointestinal tract are essential for the metabolism and absorption of nutrients and food-derived bioactive molecules. These pathways determine how these compounds are ultimately used, whether as energy sources, reaction catalysts, signaling receptor ligands, or structural components of cells [76]. Significant variation exists among individuals in the metabolic steps that govern the processing of nutrients and bioactive compounds. This leads to differences in how people respond to dietary components, and a better understanding of the factors driving this nutritional metabolic heterogeneity could help predict individual responses more accurately.

4. Nutritional Composition of Spinach

The leafy green vegetable spinach, formally known as Spinacia oleracea, is prized for its abundant nutritional value. It is a beneficial supplement to a balanced diet because it is a great source of vitamins, minerals, and antioxidants. Spinach's status as a crucial dietary component is further enhanced by the multiple health benefits it offers due to its diverse range of bioactive chemicals and metabolites [77], [78], and [79]. Its abundance of vital nutrients and bioactive substances supports many physiological processes and makes a substantial contribution to general health [80], [81], [82], [83].

The bioavailability of bioactive compounds found in foods like spinach is a complex area of research that continues to present significant challenges [84]. Bioaccessibility testing is conducted using various protocols, but currently, there is no universally accepted method for standardizing or harmonizing these procedures [85]. The efficiency of bioactive chemicals is largely determined by their bioavailability, which quantifies how well the body absorbs and uses these molecules [86]. Simply determining the concentration of bioactive compounds in food is not enough to evaluate their bioavailability or bioaccessibility. This is because the compounds that reach the bloodstream undergo complex digestion processes before exerting their effects in vivo [87].

Assessing bioaccessibility is essential for ensuring the nutritional efficacy of food products. It provides important information that helps in selecting the appropriate dosage and identifying the best food sources [88]. However, nutritional and environmental metrics should be interpreted cautiously, considering the variability introduced by factors like food biochemistry and individual human physiology [88], [89].

The bioaccessibility and bioavailability of nutrients and bioactive substances are assessed using both in vitro and in vivo techniques. In vitro assessments offer an important preliminary screening tool, but in vivo studies are still considered the gold standard for accurately evaluating bioactive compound absorption and utilization within specific populations [90]. Several factors, including the food matrix, processing methods, and individual physiological traits, play major roles in influencing the bioaccessibility and bioavailability of spinach components [85].

Water spinach, in particular, has gained scientific interest for its abundance of minerals, vitamins, antioxidants, and other bioactive substances [1]. Studies have shown that the bioaccessibility of chromium in water spinach is higher compared to other leafy vegetables like leaf lettuce and celery, making it a particularly valuable nutritional resource [91].

5. Health Benefits of Spinach

Due to its distinct nutrient and bioactive chemical content, spinach consumption is linked to a number of health benefits.

5.1 Cardiovascular Health

Nitrates, which are abundant in spinach, are transformed by the body into nitric oxide, a substance that is known to enhance blood flow and encourage vasodilation. Consuming spinach on a regular basis may help lower blood pressure and lower the risk of cardiovascular illnesses.

5.2 Cancer Prevention

Numerous antioxidants included in spinach, such as vitamins C and E, flavonoids, and carotenoids, help shield cells from harm brought on by free radicals. By neutralizing free radicals, these antioxidants help to lessen oxidative stress and the risk of developing chronic illnesses like cancer. Consuming spinach on a regular basis may help prevent cancer cells from growing and spreading [92].

Health benefits derived from cruciferous foods are closely associated with their bioactive nutrients and phytochemical compounds, particularly phenols [93]. Similarly, the leaves of African nightshade are rich in essential nutrients, including proteins, iron, ascorbic acid, and riboflavin, making it a vital nutritional source, especially for individuals from lower economic backgrounds [4].

5.3 Anti-inflammatory and Antioxidant Properties

The flavonoids, carotenoids, and phenolic compounds found in spinach have potent antioxidant and anti-inflammatory qualities. These compounds can help prevent diseases like cancer, heart disease, and neurological problems that are linked to oxidative stress by lowering inflammation and scavenging free radicals [94].

5.2 Cardiovascular Health

The nitrate in spinach is essential for lowering blood pressure because it causes vasodilation. Additionally, by balancing electrolyte levels and promoting regular muscle activity, spinach's high potassium and magnesium content helps to maintain a healthy heart and circulatory system [95].

5.3 Diabetes and Metabolic Syndrome

Because of its high fiber content and low glycemic index, spinach has demonstrated promise in controlling blood sugar levels. Spinach's bioactive ingredients may also increase insulin sensitivity, which is important for both managing and preventing type 2 diabetes [96].

5.4 Bone Health

The presence of vitamin K1, calcium, and magnesium in spinach supports bone mineralization and maintenance. Because vitamin K1 is essential for calcium metabolism and bone protein production, spinach is a good diet for healthy bones, especially in older persons [94]

5.5 Eye Health

The high carotenoid content of spinach, especially lutein and zeaxanthin, promotes eye health by shielding the retina from oxidative stress and blue light damage. These carotenoids have been associated with a lower risk of cataracts and age-related macular degeneration [94].

5.6 Cognitive and Neurological Benefits

By lowering oxidative stress and inflammation in the brain, spinach's antioxidants may help prevent cognitive loss. Studies suggest that regular spinach consumption may have neuroprotective effects, contributing to improved memory and cognitive function.[94]

5.7 Digestive Health

Spinach’s fiber content is beneficial for digestion. It enhances overall gastrointestinal health, lowers the risk of constipation, and helps regulate bowel motions. Additionally, spinach's high water content aids in maintaining hydration and digestive efficiency.[94]

6. Potential Risks and Limitations

6.1 Oxalate Content

Oxalates, which are found in spinach, have the ability to bind to calcium and other minerals, potentially reducing their absorption. Individuals who are prone to kidney stones or suffer from mineral deficiencies are advised to consume spinach in moderation. Cooking spinach can help lower its oxalate levels and may make mineral absorption more efficient. When compared to other leafy green vegetables, spinach intake is still low despite its many beneficial qualities [97]. Factors such as taste preferences, availability, and cultural eating habits greatly influence how much spinach people include in their diets. Dietary interventions using spinach have shown promise in improving insulin resistance, endothelial function, and inflammation markers in mice [98]. However, the bioavailability of nutrients derived from spinach remains an important factor to consider.

7. Impact of Processing and Cooking on Nutritional Value

Processing and cooking methods are essential in forming the nutritional composition of spinach, directly impacting how readily its nutrients and bioactive substances are absorbed. Different culinary treatments bring about varying effects on spinach's nutritional profile. For example, blanching, steaming, and sautéing help enhance the bioavailability of carotenoids and flavonoids by dissolving the cell walls and releasing these compounds from their bound forms within the plant's cellular matrix [97]. However, prolonged boiling can cause a significant loss of water-soluble vitamins, including folate and vitamin C, as these nutrients tend to leach into the cooking water. Steaming is often seen as one of the greatest ways to maintain spinach's nutritional value, since it minimizes nutrient loss while boosting the bioavailability of certain beneficial compounds. Methods such as stir-frying over high heat with a tiny amount of oil can also help retain much of spinach's nutritional value, provided the cooking time is kept short to avoid excessive degradation. On the other hand, methods such as deep-frying or prolonged boiling can greatly reduce the levels of important nutrients. The extent of nutrient degradation depends on factors like temperature, cooking duration, and the amount of water used during the cooking process [99].The nutrient retention in cooked spinach is shown in Figure 2.



Figure 2 Nutrition retention in cooked spinach

8. Future Prospects and Research Directions

Future studies should focus on methods to increase the minerals and bioactive chemicals in spinach's bioavailability. Exploring novel food processing techniques, such as pulsed electric field processing and enzymatic treatments, holds promise for enhancing nutrient extraction and retention, thereby maximizing the advantages of spinach eating for health [100]. Additional studies are needed to better understand the synergistic effects between spinach's various bioactive compounds and their interactions with other components in the diet. Clinical trials are also crucial to confirm the health benefits observed in preclinical studies and to develop optimal consumption guidelines tailored to different population groups. Different parts of water spinach, including the leaves, petioles, and stems, show notable differences in nutritional composition; stems generally contain higher concentrations of crude protein, fiber, and some fat, whereas the leaves are richer in carbohydrates and tend to have an alkaline character [1]. Research is also systematically evaluating the effects of spinach extract on physical performance, with the expectation that it may enhance performance with minimal side effects [101]. Furthermore, spinach consumption has been linked to improved cardiovascular health, likely linked to the vasoprotective effects of nitric oxide produced from the vegetable's high nitrate content [102]. In healthy people, eating foods high in nitrates, such as spinach, has been demonstrated to lower blood pressure and arterial stiffness [103].

CONCLUSION

Spinach stands out as an incredibly nutritious and health-promoting vegetable. Its high levels of essential nutrients and powerful bioactive compounds make it a smart choice for supporting overall health. While its oxalate content can slightly limit mineral absorption, this does not overshadow its many benefits. With improvements in farming techniques and food processing technologies, there is great potential to boost the nutritional impact of spinach even further. Moving forward, research focused on improving nutrient bioavailability, refining bioaccessibility testing methods, and validating health claims through clinical studies will be key. By better understanding and enhancing spinach’s full potential, we can ensure it remains a vital part of healthy diets and sustainable food systems.

REFERENCES

[1] A. H. M. Shariff et al., "Proximate compositions of Ipomea aquatic Forsk. (leaf, petiole and stem) from LubokBungor, Jeli, Kelantan," AIP conference proceedings, Jan. 2019, doi: 10.1063/1.5125564.

[2] H. Hasanah, E. Pangestu, A. Agus, and J. Achmadi, "In vitro Rumen Fermentability of the Pelleted Feed Containing Water Spinach (Ipomoea aquatica) ," American Journal of Animal and Veterinary Sciences, vol. 15, no. 1, p. 67, Jan. 2020, doi: 10.3844/ajavsp.2020.67.72.

[3] A. Gupta, N. Yadav, and S. Verma, "Nutritional evaluation of value added products incorporated with underutilized water spinach (Ipomoea aquatica) leaves," ActaHorticulturae, no. 1241, p. 549, Jun. 2019, doi: 10.17660/actahortic.2019.1241.81.

[4] G. A. Odongo et al., "African Nightshade (Solanumscabrum Mill.): Impact of Cultivation and Plant Processing on Its Health Promoting Potential as Determined in a Human Liver Cell Model," Nutrients, vol. 10, no. 10, p. 1532, Oct. 2018, doi: 10.3390/nu10101532.

[5] M. Chepkirui, P. Orina, M. A. Opiyo, P. Muendo, K. Mbogo, and R. Omondi, "Growth performance of Nile Tilapia (Oreochromisniloticus) fingerlings fed with water spinach (Ipomoea aquatica) diets," Annals of Marine Science, vol. 6, no. 1, p. 1, Apr. 2022, doi: 10.17352/ams.000026.

[6] H. M. Amir et al., "Macronutrient Concentration in Stem, Leaf and Petiole of Wild Grown Water Spinach (Ipomea Aquatic Forsk.) and Its Relationship With Pond Water," IOP Conference Series Earth and Environmental Science, vol. 709, no. 1, p. 12080, Mar. 2021, doi: 10.1088/1755-1315/709/1/012080.

[7] B. Lakitan and K. Kartika, "Population density, multiple harvesting, and ability of Ipomoea reptans to compete with native weeds at tropical wetlands," Biodiversitas Journal of Biological Diversity, vol. 21, no. 9, Aug. 2020, doi: 10.13057/biodiv/d210957.

[8] Y. Andriani, Y. Dhahiyat, Z. Zahidah, and I. Zidni, "The effect of stocking density ratio of fish on water plant productivity in aquaponics culture system," Nusantara Bioscience, vol. 9, no. 1, p. 31, Jan. 2017, doi: 10.13057/nusbiosci/n090106.

[9] Y. Andriani, Z. Zahidah, Y. Dhahiyat, H. Hamdani, and D. R. Dewi, "Performance of Lettuce and Water Spinach in Koi Fish-based Aquaponics System," Asian Journal of Fisheries and Aquatic Research, p. 1, Jul. 2019, doi: 10.9734/ajfar/2019/v3i430039.

[10] T. P. Quí, A. Ardi, and I. Chaniago, "Compare the growth and productivity of I. aquatic species on hydroponic subsystems within an aquaponic system," IOP Conference Series Earth and Environmental Science, vol. 497, no. 1, p. 12004, Apr. 2020, doi: 10.1088/1755-1315/497/1/012004.

[11] C. V. Nanda and M. N. Khozin, "Cultivation of Water Spinach Using A Hydroponic Systems at Different AB Mix Concentrations," Journal of Soilscape and Agriculture, vol. 1, no. 1, p. 1, Sep. 2022, doi: 10.19184/jsa.v1i1.125.

[12] G. Zipporah, "The Potential of Aquaponics as Food Production and Nutrient Recovery Systems in Kenya," International Scientific Days 2018. Towards Productive, Sustainable and Resilient Global Agriculture and Food Systems: Proceedings, p. 1154, Dec. 2018, doi: 10.15414/isd2018.s4.21.

[13] A. Endut, F. Lananan, S. H. A. Hamid, A. Jusoh, and W. B. W. Nik, "Balancing of nutrient uptake by water spinach (Ipomoea aquatica) and mustard green (Brassica juncea) with nutrient production by African catfish (Clariasgariepinus) in scaling aquaponic recirculation system," Desalination and Water Treatment, vol. 57, no. 60, p. 29531, May 2016, doi: 10.1080/19443994.2016.1184593.

[14] R. M. Thomas, A. K. Verma, C. Prakash, H. Krishna, S. Prakash, and A. Kumar, "Utilization of Inland saline underground water for bio-integration of Nile tilapia (Oreochromisniloticus) and spinach (Spinaciaoleracea)," Agricultural Water Management, vol. 222, p. 154, Jun. 2019, doi: 10.1016/j.agwat.2019.06.001.

[15] M. Nichols and N. Savidov, "AQUAPONICS: PROTEIN AND VEGETABLES FOR DEVELOPING COUNTRIES," ActaHorticulturae, no. 958, p. 189, Aug. 2012, doi: 10.17660/actahortic.2012.958.22.

[16] M. S. Hussein, A. ELzayat, and M. AdelBauomi, "Study of the effects of different monosex Nile tilapia (Oreochromisniloticus) fingerlings densitieson Lettuce (Lactucasativa) and water quality in a low-techrecirculationaquaponic system," International Journal of Development, vol. 9, no. 1, p. 11, Mar. 2020, doi: 10.21608/idj.2020.74591.

[17] O. Askari-Khorasgani and M. Pessarakli, "Tomato (Solanumlycopersicum) culture in vermi-aquaponic systems: II. Strategies for sustainable and economic development: Fertilization practices in vermi-ponic unit," Journal of Plant Nutrition, vol. 43, no. 11, p. 1726, Mar. 2020, doi: 10.1080/01904167.2020.1739307.
[18] P. Stathopoulou et al., "Basil and Nile tilapia Production in a Small Scale Aquaponic System," Journal of FisheriesSciences com, vol. 12, no. 4, Jan. 2018, doi: 10.21767/1307-234x.1000152.
[19] R. Ghamkhar, C. Hartleb, F. Wu, and A. Hicks, "Life cycle assessment of a cold weather aquaponic food production system," Journal of Cleaner Production, vol. 244, p. 118767, Oct. 2019, doi: 10.1016/j.jclepro.2019.118767.
[20] M. Shafahi and D. Woolston, "Aquaponics: A Sustainable Food Production System," Nov. 2014, doi: 10.1115/imece2014-39441.
[21] R. Nelson, "Aquaponics." p. 246, Dec. 21, 2016. doi: 10.1002/9781118970652.ch15.
[22] D. K. Binawati and D. A. Puspita, "THE GROWTH AND YIELDS OF LETTUCE (LACTUCA SATIVA L.) ON A AQUAPONIC NFT (NUTRIENT FILM TECHNIQUE) SYSTEM WITH THREE DIFFERENT TYPES OF FISH," Journal of Scientific Health, p. 85, Apr. 2023, doi: 10.56943/jsh.v2i2.293.
[23] V. T. Okomoda, S. A. Oladimeji, S. G. Solomon, S. O. Olufeagba, S. I. Ogah, and M. Ikhwanuddin, "Aquaponics production system: A review of historical perspective, opportunities, and challenges of its adoption," Food Science & Nutrition, vol. 11, no. 3. Wiley, p. 1157, Dec. 18, 2022. doi: 10.1002/fsn3.3154.
[24] A. A. Kulkarni, P. Dhanush, B. S. Chethan, T. G. C.S., and P. K. Shrivastava, "A Brief Study on Aquaponics: An Innovative Farming Technology," Indian Journal of Science and Technology, vol. 12, no. 48, p. 1, Dec. 2019, doi: 10.17485/ijst/2019/v12i48/149387.
[25] B. Himabindhu, S. Divakar. Dr. S. P. P, B. Joseph, and K. U, "Quality evaluation of water spinach (Ipomeaaquatica) leaves cultivated through Aquaponics," International Journal of Chemical Studies, vol. 9, no. 1, p. 1245, Jan. 2021, doi: 10.22271/chemi.2021.v9.i1r.11393.
[26] F. P. Alloggia, R. F. Bafumo, D. Ramírez, M. Maza, and A. B. Camargo, "Brassicaceaemicrogreens: A novel and promissory source of sustainable bioactive compounds," Current Research in Food Science, vol. 6. Elsevier BV, p. 100480, Jan. 01, 2023. doi: 10.1016/j.crfs.2023.100480.
[27] N. Xi, Y. Li, J. Chen, Y. Yixiao, J. Duan, and X. Xia, "Elevated Temperatures Decrease the Photodegradation Rate of Pyrethroid Insecticides on Spinach Leaves: Implications for the Effect of Climate Warming," Environmental Science & Technology, vol. 55, no. 2, p. 1167, Dec. 2020, doi: 10.1021/acs.est.0c06959.
[28] S. Kaur, N. Trehan, and B. Kaur, "Analysis of spinach (Spinaciaoleracea) pigments using chromatography and UV-visible spectroscopy.," Agriways, vol. 7, no. 2, p. 96, Jan. 2019, Accessed: Jan. 2025. [Online]. Available: https://www.cabdirect.org/cabdirect/abstract/20203185527?q=(similar%3a20113026453)
[29] A. Fratianni, A. D'Agostino, S. Niro, A. Bufano, B. Paura, and G. Panfili, "Loss or Gain of Lipophilic Bioactive Compounds in Vegetables after Domestic Cooking? Effect of Steaming and Boiling," Foods, vol. 10, no. 5, p. 960, Apr. 2021, doi: 10.3390/foods10050960.
[30] Y. Li et al., "Far-red light suppresses glucosinolate profiles of Chinese kale through inhibiting genes related to glucosinolate biosynthesis," Environmental and Experimental Botany, vol. 188, p. 104507, May 2021, doi: 10.1016/j.envexpbot.2021.104507.
[31] P. Lengkidworraphiphat, R. Wongpoomchai, S. Taya, and S. Jaturasitha, "Effect of genotypes on macronutrients and antioxidant capacity of chicken breast meat," Asian-Australasian Journal of Animal Sciences, vol. 33, no. 11, p. 1817, Jan. 2020, doi: 10.5713/ajas.19.0736.
[32] M. Simsek and K. Whitney, "Examination of Primary and Secondary Metabolites Associated with a Plant-Based Diet and Their Impact on Human Health," Foods, vol. 13, no. 7, p. 1020, Mar. 2024, doi: 10.3390/foods13071020.

[33] A. Valdés-Jiménez et al., "PSC-db: A Structured and Searchable 3D-Database for Plant Secondary Compounds," Molecules, vol. 26, no. 4, p. 1124, Feb. 2021, doi: 10.3390/molecules26041124.
[34] D. L. Luthria, "PHENOLIC COMPOUNDS ANALYSIS IN FOODS AND DIETARY SUPPLEMENTS IS NOT THE SAME USING DIFFERENT SAMPLE PREPARATION PROCEDURES," ActaHorticulturae, no. 841, p. 381, Aug. 2009, doi: 10.17660/actahortic.2009.841.48.
[35] E. N. Gomes et al., "Successive harvests affect the aromatic and polyphenol profiles of novel catnip (Nepetacataria L.) cultivars in a genotype-dependent manner," Frontiers in Plant Science, vol. 14, Feb. 2023, doi: 10.3389/fpls.2023.1121582.
[36] V. K. Srivastav, C. Egbuna, and M. Tiwari, "Plant secondary metabolites as lead compounds for the production of potent drugs," in Elsevier eBooks, Elsevier BV, 2020, p. 3.doi: 10.1016/b978-0-12-817890-4.00001-9.
[37] Z. A. Reshi, W. Ahmad, A. C. Лукаткин, and S. B. Javed, "From Nature to Lab: A Review of Secondary Metabolite Biosynthetic Pathways, Environmental Influences, and In Vitro Approaches," Metabolites, vol. 13, no. 8. Multidisciplinary Digital Publishing Institute, p. 895, Jul. 28, 2023. doi: 10.3390/metabo13080895.
[38] T. Isah, "Stress and defense responses in plant secondary metabolites production," Biological Research, vol. 52, no. 1.BioMed Central, Jul. 29, 2019. doi: 10.1186/s40659-019-0246-3.
[39] A. Khan et al., "Plant Secondary Metabolites—Central Regulators Against Abiotic and Biotic Stresses," Feb. 2025, doi: 10.20944/preprints202502.0739.v1.
[40] A. D. Kinghorn, "The Discovery of Drugs from Higher Plants," Elsevier eBooks. Elsevier BV, p. 81, Jan. 01, 1994. doi: 10.1016/b978-0-7506-9003-4.50010-1.
[41] N. Khan and S. Dhawan, "Role of Molecular Markers in Assessing Genetic Diversity in Mentha: A Review," Scientific Journal of Genetics and Gene Therapy, vol. 2, no. 1. p. 22, Dec. 30, 2016. doi: 10.17352/sjggt.000011.
[42] A. Upadhyay, I. Upadhyaya, A. K. Johny, and K. Venkitanarayanan, "Combating Pathogenic Microorganisms Using Plant-Derived Antimicrobials: A Minireview of the Mechanistic Basis," BioMed Research International, vol. 2014. Hindawi Publishing Corporation, p. 1, Jan. 01, 2014. doi: 10.1155/2014/761741.
[43] A. M. L. Seca and A. Трендафилова, "Secondary Metabolites in Edible Species: Looking beyond Nutritional Value," Foods, vol. 10, no. 5, p. 1131, May 2021, doi: 10.3390/foods10051131.
[44] A. Ullah, A. Bano, and H. T. Janjua, "Microbial Secondary Metabolites and Defense of Plant Stress," in Elsevier eBooks, Elsevier BV, 2020, p. 37.doi: 10.1016/b978-0-12-819978-7.00003-8.
[45] D. Thirumurugan, A. Cholarajan, S. Raja, and R. Vijayakumar, "An Introductory Chapter: Secondary Metabolites," in InTech eBooks, 2018. doi: 10.5772/intechopen.79766.
[46] S. L. Kochhar and S. K. Gujral, "Secondary Plant Metabolites," PLANT PHYSIOLOGY. Oxford University Press, p. 590, May 15, 2020. doi: 10.1017/9781108486392.022.
[47] A. Dekebo, "Introductory Chapter: Plant Extracts," in IntechOpen eBooks, IntechOpen, 2019. doi: 10.5772/intechopen.85493.
[48] M. A. Bernards, "Plant natural products: a primerThe present review is one in the special series of reviews on animal–plant interactions.," Canadian Journal of Zoology, vol. 88, no. 7, p. 601, Jul. 2010, doi: 10.1139/z10-035.
[49] C. Troncoso et al., "CHEMICAL DEFENSE RESPONSES IN Eucalyptus globulus (Labill) PLANTS," Journal of the Chilean Chemical Society, vol. 56, no. 3, p. 768, Jan. 2011, doi: 10.4067/s0717-97072011000300010.
[50] A. Plášková and J. Mlček, "New insights of the application of water or ethanol-water plant extract rich in active compounds in food," Frontiers in Nutrition, vol. 10. Frontiers Media, Mar. 28, 2023. doi: 10.3389/fnut.2023.1118761.
[51] Y. Li, K. Wen, X. Ruan, Y. Zhao, W. Feng, and Q. Wang, "Response of Plant Secondary Metabolites to Environmental Factors," Molecules, vol. 23, no. 4. Multidisciplinary Digital Publishing Institute, p. 762, Mar. 27, 2018. doi: 10.3390/molecules23040762.

[52] A. Armarkar, D. S. Mahure, A. R. Pounikar, R. T. Bhagat, and D. N. Nagare, "An Overview on the Biosynthetic Pathways and Medicinal Values of Secondary Metabolites," Journal of Pharmaceutical Research International, p. 100, Jun. 2021, doi: 10.9734/jpri/2021/v33i33a31777.
[53] J. S. Câmara, R. Perestrelo, R. Ferreira, C. V. Berenguer, J. A. M. Pereira, and P. C. Castilho, "Plant-Derived Terpenoids: A Plethora of Bioactive Compounds with Several Health Functions and Industrial Applications—A Comprehensive Overview," Molecules, vol. 29, no. 16. Multidisciplinary Digital Publishing Institute, p. 3861, Aug. 15, 2024. doi: 10.3390/molecules29163861.
[54] A. B. Abtew, S. Subramanian, X. Cheseto, S. Kreiter, G. T. Garzia, and T. Martin, "Repellency of Plant Extracts against the Legume Flower ThripsMegalurothripssjostedti (Thysanoptera: Thripidae)," Insects, vol. 6, no. 3, p. 608, Jun. 2015, doi: 10.3390/insects6030608.
[55] M. Wink, "Potential of DNA Intercalating Alkaloids and Other Plant Secondary Metabolites against SARS-CoV-2 Causing COVID-19," Diversity, vol. 12, no. 5, p. 175, Apr. 2020, doi: 10.3390/d12050175.
[56] H. Sánchez-Sánchez and A. Morquecho-Contreras, "Chemical Plant DefenseAgainst Herbivores," in InTech eBooks, 2017. doi: 10.5772/67346.
[57] K. Edwina and P. Leela, "Phytotoxic Effect OfCyanodonDactylon (L.) Pers. and CyperusRotundus L. On Growth And Biochemical Changes of VignaRadiata (L.)R. Wilczek," International Research Journal on Advanced Science Hub, vol. 2, p. 90, Nov. 2020, doi: 10.47392/irjash.2020.166.
[58] Md. A. Rahman, T. Ahsan, and S. Islam, "Antibacterial and antifungal properties of the methanol extract from the stem of Argyreiaargentea," Bangladesh Journal of Pharmacology, vol. 5, no. 1, Mar. 2010, doi: 10.3329/bjp.v5i1.4700.
[59] P. A. Divekar et al., "Plant Secondary Metabolites as Defense Tools against Herbivores for Sustainable Crop Protection," International Journal of Molecular Sciences, vol. 23, no. 5. Multidisciplinary Digital Publishing Institute, p. 2690, Feb. 28, 2022. doi: 10.3390/ijms23052690.
[60] A. R. War et al., "Mechanisms of plant defense against insect herbivores," Plant Signaling&Behavior, vol. 7, no. 10. Taylor & Francis, p. 1306, Sep. 30, 2012. doi: 10.4161/psb.21663.
[61] J. P. Yactayo-Chang, H. V. Tang, J. Mendoza, S. A. Christensen, and A. K. Block, "Plant Defense Chemicals against Insect Pests," Agronomy, vol. 10, no. 8, p. 1156, Aug. 2020, doi: 10.3390/agronomy10081156.
[62] M. I. Haggag, "Phytochemical profile for Cestrum nocturnum leaves ethanolic extract and isolation of a rare flavonoid using different chromatographic and spectroscopic techniques," Journal of Medicinal Plants Studies, vol. 10, no. 2, p. 143, Mar. 2022, doi: 10.22271/plants.2022.v10.i2b.1403.
[63] D. Balmer, V. Flors, G. Glauser, and B. Mauch-Mani, "Metabolomics of cereals under biotic stress: current knowledge and techniques," Frontiers in Plant Science, vol. 4, Jan. 2013, doi: 10.3389/fpls.2013.00082.
[64] M. D. Bolton, "Primary Metabolism and Plant Defense—Fuel for the Fire," Molecular Plant-Microbe Interactions, vol. 22, no. 5. American Phytopathological Society, p. 487, Apr. 06, 2009. doi: 10.1094/mpmi-22-5-0487.
[65] U. Salam et al., "Plant Metabolomics: An Overview of the Role of Primary and Secondary Metabolites against Different Environmental Stress Factors," Life, vol. 13, no. 3. Multidisciplinary Digital Publishing Institute, p. 706, Mar. 06, 2023. doi: 10.3390/life13030706.
[66] S. Rasmann and A. A. Agrawal, "Plant defense against herbivory: progress in identifying synergism, redundancy, and antagonism between resistance traits," Current Opinion in Plant Biology, vol. 12, no. 4. Elsevier BV, p. 473, Jun. 19, 2009. doi: 10.1016/j.pbi.2009.05.005.
[67] S. Selma, N. Ntelkis, T. H. Nguyen, and A. Goossens, "Engineering the plant metabolic system by exploiting metabolic regulation," The Plant Journal, vol. 114, no. 5, p. 1149, Feb. 2023, doi: 10.1111/tpj.16157.
[68] G. Niu and H. Tan, "Biosynthesis and regulation of secondary metabolites in microorganisms," Science China Life Sciences, vol. 56, no. 7, p. 581, Jul. 2013, doi: 10.1007/s11427-013-4501-5.

[69] C. M. Rojas, M. Senthil-Kumar, V. Tzin, and K. S. Mysore, "Regulation of primary plant metabolism during plant-pathogen interactions and its contribution to plant defense," Frontiers in Plant Science, vol. 5. Frontiers Media, Jan. 01, 2014. doi: 10.3389/fpls.2014.00017.
[70] C. Zhan, "Microbial Decomposition and Soil Health: Mechanisms and Ecological Implications," Molecular Soil Biology, Jan. 2024, doi: 10.5376/msb.2024.15.0007.
[71] F. Yang, X. Ding, and G. Lv, "Quantitative proteomic analysis based on TMT reveals different responses of Haloxylonammodendron and Haloxylonpersicum to long-term drought," BMC Plant Biology, vol. 25, no. 1, Apr. 2025, doi: 10.1186/s12870-025-06513-x.
[72] K. Kytidou, M. Artola, H. S. Overkleeft, and J. M. F. G. Aerts, "Plant Glycosides and Glycosidases: A Treasure-Trove for Therapeutics," Frontiers in Plant Science, vol. 11. Frontiers Media, Apr. 07, 2020. doi: 10.3389/fpls.2020.00357.
[73] A. M. Boutanaev et al., "Investigation of terpene diversification across multiple sequenced plant genomes," Proceedings of the National Academy of Sciences, vol. 112, no. 1, Dec. 2014, doi: 10.1073/pnas.1419547112.
[74] S. Oh et al., "Approach strategies and application of metabolomics to biotechnology in plants," Frontiers in Plant Science, vol. 14. Frontiers Media, Aug. 11, 2023. doi: 10.3389/fpls.2023.1192235.
[75] F. J. Barba, M. J. Esteve, and A. Frígola, "Bioactive Components from Leaf Vegetable Products," in Studies in natural products chemistry, Elsevier BV, 2014, p. 321. doi: 10.1016/b978-0-444-63294-4.00011-5.
[76] S. H. Zeisel, "Precision (Personalized) Nutrition: Understanding Metabolic Heterogeneity," Annual Review of Food Science and Technology, vol. 11, no. 1. Annual Reviews, p. 71, Jan. 13, 2020. doi: 10.1146/annurev-food-032519-051736.
[77] F. Zhu, W. Wen, Y. Cheng, S. Alseekh, and A. R. Fernie, "Integrating multiomics data accelerates elucidation of plant primary and secondary metabolic pathways," aBIOTECH, vol. 4, no. 1. Springer Science+Business Media, p. 47, Jan. 11, 2023. doi: 10.1007/s42994-022-00091-4.
[78] K. Wojtunik-Kulesza, A. Oniszczuk, T. Oniszczuk, M. Combrzyński, D. Nowakowska, and A. Matwijczuk, "Influence of In Vitro Digestion on Composition, Bioaccessibility and Antioxidant Activity of Food Polyphenols—A Non-Systematic Review," Nutrients, vol. 12, no. 5. Multidisciplinary Digital Publishing Institute, p. 1401, May 13, 2020. doi: 10.3390/nu12051401.
[79] R. A. Hussein and A. A. El-Anssary, "Plants Secondary Metabolites: The Key Drivers of the Pharmacological Actions of Medicinal Plants," in IntechOpen eBooks, IntechOpen, 2019. doi: 10.5772/intechopen.76139.
[80] M. Kussmann, D. H. A. Cunha, and S. Berciano, "Bioactive compounds for human and planetary health," Frontiers in Nutrition, vol. 10. Frontiers Media, Jul. 17, 2023. doi: 10.3389/fnut.2023.1193848.
[81] C. O. Ajanaku et al., "Functional bioactive compounds in ginger, turmeric, and garlic," Frontiers in Nutrition, vol. 9. Frontiers Media, Dec. 08, 2022. doi: 10.3389/fnut.2022.1012023.
[82] S. Noore, N. K. Rastogi, C. P. O'Donnell, and B. K. Tiwari, "Novel Bioactive Extraction and Nano-Encapsulation," Encyclopedia, vol. 1, no. 3, p. 632, Jul. 2021, doi: 10.3390/encyclopedia1030052.
[83] A. Boronat et al., "Contribution of Biotransformations Carried Out by the Microbiota, Drug-Metabolizing Enzymes, and Transport Proteins to the Biological Activities of Phytochemicals Found in the Diet," Advances in Nutrition, vol. 12, no. 6. Elsevier BV, p. 2172, Aug. 09, 2021. doi: 10.1093/advances/nmab085.
[84] M. J. Rein, M. Renouf, C. Cruz-Hernandez, L. Actis-Goretta, S. K. Thakkar, and M. da S. Pinto, "Bioavailability of bioactive food compounds: a challenging journey to bioefficacy," British Journal of Clinical Pharmacology, vol. 75, no. 3. Wiley, p. 588, Aug. 16, 2012. doi: 10.1111/j.1365-2125.2012.04425.x.
[85] F. Cámara-Martos, S. Obregón, O. Mesa-Plata, M. E. Cartea, and A. de H. Bailón, "Quantification and in vitro bioaccessibility of glucosinolates and trace elements in Brassicaceae leafy vegetables," Food Chemistry, vol. 339, p. 127860, Aug. 2020, doi: 10.1016/j.foodchem.2020.127860.

[86] M. D'Archivio, C. Filesi, R. Varí, B. Scazzocchio, and R. Masella, "Bioavailability of the Polyphenols: Status and Controversies," International Journal of Molecular Sciences, vol. 11, no. 4. Multidisciplinary Digital Publishing Institute, p. 1321, Mar. 31, 2010. doi: 10.3390/ijms11041321.
[87] D. I. Santos, J. A. Saraiva, A. A. Vicente, and M. Moldão-Martins, "Methods for determining bioavailability and bioaccessibility of bioactive compounds and nutrients," in Elsevier eBooks, Elsevier BV, 2019, p. 23. doi: 10.1016/b978-0-12-814174-8.00002-0.
[88] J. M. Carbonell-Capella, M. Buniowska, F. J. Barba, M. J. Esteve, and A. Frígola, "Analytical Methods for Determining Bioavailability and Bioaccessibility of Bioactive Compounds from Fruits and Vegetables: A Review," Comprehensive Reviews in Food Science and Food Safety, vol. 13, no. 2. Wiley, p. 155, Feb. 21, 2014. doi: 10.1111/1541-4337.12049.
[89] F. Leroy, T. Beal, P. Gregorini, G. A. McAuliffe, and S. van Vliet, "Nutritionism in a food policy context: the case of 'animal protein,'" Animal Production Science, vol. 62, no. 8, p. 712, Feb. 2022, doi: 10.1071/an21237.
[90] E. Fernández-García, I. Carvajal-Lérida, and A. Pérez-Gálvez, "In vitro bioaccessibility assessment as a prediction tool of nutritional efficiency," Nutrition Research, vol. 29, no. 11. Elsevier BV, p. 751, Nov. 01, 2009. doi: 10.1016/j.nutres.2009.09.016.
[91] P. Wang et al., "Variability of chromium bioaccessibility and speciation in vegetables: The influence of in vitro methods, gut microbiota and vegetable species," Food Chemistry, vol. 277, p. 347, Oct. 2018, doi: 10.1016/j.foodchem.2018.10.120.
[92] N. Satheesh and S. W. Fanta, "Kale: Review on nutritional composition, bio-active compounds, anti-nutritional factors, health beneficial properties and value-added products," Cogent Food & Agriculture, vol. 6, no. 1, p. 1811048, Jan. 2020, doi: 10.1080/23311932.2020.1811048.
[93] A. Costa-Pérez et al., "Systematic Review on the Metabolic Interest of Glucosinolates and Their Bioactive Derivatives for Human Health," Nutrients, vol. 15, no. 6. Multidisciplinary Digital Publishing Institute, p. 1424, Mar. 15, 2023. doi: 10.3390/nu15061424.
[94] C. S. Lasya, "Spinach and its health benefits: A review." Aug. 2022. Accessed: Apr. 27, 2025.[Online]. Available: https://www.thepharmajournal.com/archives/2022/vol11issue8S/PartP/S-11-8-47-875.pdf
[95] J. Singh, G. K. Jayaprakasha, and B. S. Patil, "Extraction, Identification, and Potential Health Benefits of Spinach Flavonoids: A Review," ACS symposium series. American Chemical Society, p. 107, Jan. 01, 2018. doi: 10.1021/bk-2018-1286.ch006.
[96] P. Mirmiran, "Functional foods-based diet as a novel dietary approach for management of type 2 diabetes and its complications: A review," World Journal of Diabetes, vol. 5, no. 3. p. 267, Jan. 01, 2014. doi: 10.4239/wjd.v5.i3.267.
[97] J. L. Roberts and R. Moreau, "Functional properties of spinach (Spinaciaoleracea L.) phytochemicals and bioactives," Food & Function, vol. 7, no. 8. Royal Society of Chemistry, p. 3337, Jan. 01, 2016. doi: 10.1039/c6fo00051g.
[98] T. Li, X. Lu, Y. Sun, and X. Yang, "Effects of spinach nitrate on insulin resistance, endothelial dysfunction markers and inflammation in mice with high-fat and high-fructose consumption," Food & Nutrition Research, vol. 60, no. 1, p. 32010, Jan. 2016, doi: 10.3402/fnr.v60.32010.
[99] A. V. Barker and D. N. Maynard, "Nutritional factors affecting nitrate accumulation in spinach," Communications in Soil Science and Plant Analysis, vol. 2, no. 6, p. 471, Jan. 1971, doi: 10.1080/00103627109366339.
[100] L. C. R. dos Reis, V. R. de Oliveira, M. E. K. Hagen, A. Jablonski, S. H. Flôres, and A. de O. Rios, "Carotenoids, flavonoids, chlorophylls, phenolic compounds and antioxidant activity in fresh and cooked broccoli (Brassica oleracea var. Avenger) and cauliflower (Brassica oleracea var. Alphina F1)," LWT, vol. 63, no. 1, p. 177, Apr. 2015, doi: 10.1016/j.lwt.2015.03.089.
[101] J. Wen et al., "Improved Effect of Spinach Extract on Physical Performance: A Systematic Review of Randomized Controlled Trials," Cureus. Cureus, Inc., Jan. 22, 2025. doi: 10.7759/cureus.77840.

[102] E. Jovanovski et al., "Effect of Spinach, a High Dietary Nitrate Source, on Arterial Stiffness and Related Hemodynamic Measures: A Randomized, Controlled Trial in Healthy Adults," Clinical Nutrition Research, vol. 4, no. 3, p. 160, Jan. 2015, doi: 10.7762/cnr.2015.4.3.160.

[103] A. H. Liu et al., "Effects of a nitrate-rich meal on arterial stiffness and blood pressure in healthy volunteers," Nitric Oxide, vol. 35, p. 123, Oct. 2013, doi: 10.1016/j.niox.2013.10.001.