IJCRT.ORG ISSN : 2320-2882

INTERNATIONAL JOURNAL OF CREATIVE RESEARCH THOUGHTS (IJCRT)
An International Open Access, Peer-reviewed, Refereed Journal

A Study Of Blood Alcohol Stability In Forensic Antemortem Blood Samples

[1]K R Ramya,
[1]Student,
[1]Department of Forensic Science ,
[1]Parul University , Vadodara, India

Abstract: Alcohol is extensively consumed and recognized for its significant consequences on health, regulation, and society. It is absorbed in the stomach and intestines and distributed via body fluids, typically metabolized in the liver. Initial relaxation is experienced in the brain, but higher amounts lead to drowsiness and unconsciousness. Research on its impact on young adults is crucial due to observed differences in alcohol metabolism within this age group. In forensic investigations, the stability of ethanol levels in stored blood samples has been analysed. It has been determined that factors such as temperature, preservatives, and storage conditions influence ethanol levels, potentially raising concerns about the reliability of re-examined samples in criminal proceedings. To address alcohol-related fatalities among young drivers, measures including reducing the legal drinking age and implementing "zero tolerance" rules have been successfully introduced. The detection of alcohol impairment near legal limits has been identified as challenging, with only trained experts proven to accurately understand early symptoms of intoxication. The necessity of understanding alcohol's physiological effects, legal implications, and related health risks has been emphasized to mitigate damage within society. Awareness campaigns and stringent regulations are recommended as strategies to reduce alcohol's negative societal impact.

Keywords: Metabolized, Forensic investigation, Ethanol levels, Re-tested, young drivers, zero tolerance, Blood Alcohol Concentration (BAC), Driving Under Influence (DUI).

Introduction

The stability of ethanol in antemortem blood specimens has been extensively studied due to its essential role in forensic toxicology. Pre-analytical factors, such as storage conditions, sample handling, and the use of preservatives, have been identified as key influences on ethanol concentration over time. Ethanol stability is crucial for maintaining the integrity of blood alcohol concentration (BAC) measurements, which are critical evidence in cases involving driving under the influence (DUI) [1]. This study aims to analyse how pre-analytical factors affect ethanol stability in antemortem blood samples during storage.

The analysis of blood alcohol concentration (BAC) is significantly recognized as a critical element in forensic toxicology, particularly in instances concerning driving under the influence (DUI) and other alcohol-related incidents. The use of gas chromatography-mass spectrometry (GC-MS) has been established as a reliable and sensitive method for the quantification of ethanol in blood. This method is favoured due to its high specificity, sensitivity, and potential to differentiate ethanol from specific volatile compounds. The stability of ethanol in blood samples during storage is known to be influenced by various pre-analytical factors, including temperature, preservatives, and sample handling. This study was conducted to investigate the reliability of GC-MS for BAC determination and to assess how storage conditions affected ethanol stability in forensic blood samples [2].

The stability of ethanol in antemortem blood samples saved under refrigeration for prolonged durations has been investigated. Previous studies have shown that ethanol concentration decreases over time due to factors such as microbial activity, evaporation, and oxidation. These studies aimed to assess the feasibility of measuring forensically significant blood ethanol concentrations in antemortem blood samples saved refrigerated for 4-7 years. The study was carried out to provide insights into the reliability of reanalysing blood samples for ethanol content after long-term storage [3].

The stability of ethanol in forensic blood samples has been significantly studied due to its critical importance in maintaining the reliability of blood alcohol concentration (BAC) measurements in legal and forensic investigations. Ethanol concentrations are known to decrease over time due to factors such as microbial activity, evaporation, and oxidation. Storage conditions, including temperature, length, and the type of container (glass or plastic), have been shown to seriously impact ethanol stability. This study was conducted to evaluate the effects of storage conditions on ethanol stability in blood samples stored for extended periods at refrigerated temperatures. The findings are intended to provide insights into best practices for preserving ethanol concentrations in forensic blood samples [4].

Blood alcohol concentration (BAC) is a critical piece of evidence in forensic investigations, particularly in cases involving driving under the influence (DUI). The stability of ethanol in blood samples is critical for ensuring the reliability of BAC measurements over time. Factors including storage situations, preservatives, and handling practices could significantly affect ethanol degrees, potentially impacting legal results. This study aimed to investigate the stability of blood alcohol in forensic antemortem blood samples under diverse storage conditions to provide insights into advanced storage practices for retaining sample integrity [5].

The stability of ethanol in forensic antemortem whole blood samples stored at room temperature for prolonged durations has been investigated. This study aimed to determine the effects of long-term storage on blood alcohol concentration (BAC) in preserved entire blood specimens. Ethanol stability is critical for maintaining the integrity of forensic evidence, especially in cases regarding driving under the influence (DUI). Factors such as storage situations, pattern coping with, and preservative use have been shown to influence ethanol stability, potentially impacting criminal outcomes [6].

Ethanol stability in unpreserved refrigerated antemortem blood samples was investigated to assess the reliability of blood alcohol attention (BAC) measurements over time. The absence of preservatives, inclusive of sodium fluoride, in blood samples could potentially cause ethanol degradation due to microbial hobby or oxidation. However, preceding studies had shown that ethanol did not increase in unpreserved refrigerated antemortem blood, suggesting that microbial fermentation was not a significant concern. This study aimed to assess the importance of ethanol decreases in unpreserved blood samples stored at refrigerated temperatures [7].

Method

Sample Collection and Preservation

- **Sample Selection**: Thirty-two whole blood case samples were selected for this study. Each sample consisted of two tubes of blood [4].
- **Initial Analysis**: One tube from each case was analyzed for BAC using headspace gas chromatography (HS-GC) with flame ionization detection (FID) for court proceedings related to DUI [8].
- **Storage Conditions**: All blood samples were stored under refrigeration (4°C) for a period ranging from 13 to 39 months [6].

Re-analysis and Comparison

- **Re-analysis**: After the initial storage period, both tubes of blood for each case were re-analyzed for BAC using the same HS-GC method.
- **Comparison**: The results from the re-analysis were compared to the original BAC measurements to assess changes in ethanol concentration over time [4, 7, 9, 10].

Additional Storage Conditions

- **Room Temperature Storage**: The same blood samples were then stored at room temperature for six months to evaluate the effect of warmer conditions on ethanol stability.
- **High-Temperature Storage**: Subsequently, samples were stored at 38°C for 7 and 28 days to assess the impact of elevated temperatures on BAC stability [5].

Statistical Analysis

- **Data Analysis**: Statistical analysis was performed to determine the significance of changes in BAC under different storage conditions. Mean losses in BAC were calculated for both opened and unopened tubes [11, 12].

Results

Blood samples stored at 4°C for 4-7 years exhibited minimum ethanol degradation, with mean variations in ethanol concentration recorded as -0.014 g/dL for previously opened tubes and -0.010 g/dL for unopened tubes. No increases in ethanol concentration exceeding the uncertainty of preliminary measurements were determined, and all decreases were less than 0.020 g/dL. In contrast, samples stored at room temperature for extended durations (5.4 to 10.5 years) showed significant ethanol losses, ranging from 0.005 to 0.234 g/dL, with unopened and more-than-half-full tubes demonstrating better stability. Elevated temperature storage (38°C) caused significant ethanol degradation after 28 days, while short-term storage (7 days) did not result in analytically substantial modifications. The addition of sodium fluoride (NaF) as a preservative successfully inhibited microbial activity, reducing ethanol degradation in both refrigerated and room temperature situations.

Discussion

Ethanol stability in forensic blood samples has been shown to depend extensively on storage conditions, duration, and the usage of preservatives. Refrigeration at 4°C has been identified as the most effective circumstance for maintaining ethanol concentrations over both medium-term (up to 365 days) and long-term intervals (up to 7 years). Minimal ethanol losses were determined under these conditions, with mean decreases of -0.014 g/dL for previously opened tubes and -0.010 g/dL for unopened tubes, making refrigerated storage reliable for forensic purposes.

Room temperature storage resulted in greater ethanol losses over time, with losses ranging from 0.005 to 0.234 g/dL in samples stored for 5.4-10.5 years. Tubes that were unopened and more than half full demonstrated better stability compared to opened or partially filled tubes. Elevated temperatures (38°C) brought on significant ethanol degradation after 28 days, while short-term exposure (7 days) did not result in analytically extensive changes.

The use of sodium fluoride (NaF) as a preservative was found to play a vital role in preventing microbial activity and maintaining ethanol stability. Samples stored without NaF exhibited greater ethanol degradation due to microbial activity, particularly under room temperature conditions or during prolonged storage intervals.

These findings aligned with previous studies emphasizing the importance of proper sample handling and storage protocols in forensic toxicology. Refrigeration combined with the use of preservatives ensured accurate and legally defensible blood alcohol concentration (BAC) measurements even after extended storage

Conclusion

The widespread consumption of alcohol and its significant consequences on health, regulation, and society are recognized. Its absorption, distribution, and metabolism within the body have been notably studied, with specific attention given to its effects on young adults due to variations in alcohol metabolism. The stability of ethanol levels in stored blood samples has been analysed in forensic investigations, and it has been determined that factors such as temperature, preservatives, and storage situations influenced ethanol concentrations, raising concerns about the reliability of re-tested samples in criminal cases. To reduce alcohol-related fatalities among young drivers, measures together with reducing the legal consuming age and enforcing “zero tolerance” rules have been effectively delivered. The difficulty in detecting alcohol impairment close to the legal restriction has been identified, with trained professionals shown to be the most reliable in identifying early symptoms of intoxication. The necessity of understanding alcohol’s physiological effects, legal implications, and fitness risks has been emphasized to mitigate societal harm. As a result, awareness campaigns and stringent rules have been encouraged to minimize alcohol’s negative effect on society.

References

1. Laurens JB, Sewell FJJ, Kock MM. Pre-analytical factors related to the stability of ethanol concentration during storage of ante-mortem blood alcohol specimens. J Forensic Leg Med. 2018 Aug;58:155–63.
2. Cordell RL, Pandya H, Hubbard M, Turner MA, Monks PS. GC-MS analysis of ethanol and other volatile compounds in micro-volume blood samples—quantifying neonatal exposure. Anal Bioanal Chem. 2013 May 19;405(12):4139–47.
3. Kosecki PA, Raines ME. Testing antemortem blood samples for ethanol after four to seven years of refrigerated storage. J Forensic Sci. 2022 May 28;67(3):1250–7.
4. Shajani NK, Image BA, Chu EB. The Stability of Ethanol in Stored Forensic Blood Samples. Canadian Society of Forensic Science Journal. 1989 Jan;22(4):335–9.
5. Shan X, Tiscione NB, Alford I, Yeatman DT. A study of blood alcohol stability in forensic antemortem blood samples. Forensic Sci Int. 2011 Sep;211(1–3):47–50.
6. Tiscione NB, Vacha RE, Alford I, Yeatman DT, Shan X. Long-Term Blood Alcohol Stability in Forensic Antemortem Whole Blood Samples. J Anal Toxicol. 2015 Jul;39(6):419–25.
7. Kosecki PA, Canonico E, Abbott L. Ethanol stability in unpreserved refrigerated antemortem blood. J Forensic Sci. 2023 Mar 5;68(2):682–7.
8. Martínez MA, Ballesteros S. Toxicological Findings in Two Planned Complex Suicide Cases: Ingestion of Petroleum Distillates and Subsequent Hanging. J Anal Toxicol. 2009 Jul 1;33(6):336–42.
9. G. STOJINJKOVIC MMDSSBLA. Ethanol concentration changes in blood sample during medium term refrigerated storage. 2016;20:4831–6.
10. Emel Kocak F, Ozden Isiklar O, Kocak H, Meral A. Comparison of blood ethanol stabilities in different storage periods. Biochem Med (Zagreb). 2015;57–63.
11. Olds ML. Statistical comparisons of 0.08 g/100 ml fortified blood alcohol samples from 6-ml and 10-ml gray-top tubes. J Forensic Sci. 2023 Jan 8;68(1):301–7.
12. Serpil Erdogan OEO Sen. Optimal Storage Conditions for Blood Samples, for Later Analysis in Criminal Prosecution Proceedings. J Immunol Clin Microbiol. 2018 Dec;3(4):60–7.