IJCRT.ORG ISSN : 2320-2882

INTERNATIONAL JOURNAL OF CREATIVE RESEARCH THOUGHTS (IJCRT)

An International Open Access, Peer-reviewed, Refereed Journal

Role Of Pudhina Extract In Alleviating Dysmenorrhea Among Adolescents

[1]Sneha Smrti Guha, [2]Dr Payal Sharma,

[1]Ph.D. Scholar, [2]Research Supervisor,

[1]Department of Medical Surgical Nursing, [2]Department of Mental Health Nursing,

[1]Malwanchal University, Indore, M.P. [2]Malwanchal University, Indore, M.P.

Abstract

Dysmenorrhea, defined as painful menstrual cramps, affects a large proportion of adolescent girls globally, often resulting in impaired daily functioning, reduced academic performance, and decreased quality of life. Conventional treatment typically includes non-steroidal anti-inflammatory drugs (NSAIDs) and hormonal therapies; however, concerns about adverse effects and long-term use have led to increasing interest in herbal alternatives. *Pudhina* (mint; *Mentha* spp.) extract—rich in menthol, flavonoids, and essential oils—has demonstrated analgesic, antispasmodic, and anti-inflammatory properties, which may contribute to reduced menstrual pain. This review examines the pharmacological basis for using pudhina extract, evaluates clinical evidence of its effectiveness in dysmenorrhea, and discusses practical implications for adolescent health. While current studies suggest that mint extract may decrease pain intensity and associated symptoms, further high-quality randomized controlled trials in adolescent cohorts are recommended to validate these findings and establish standardized formulations.

Keywords: Dysmenorrhea, adolescents, pudhina.

1. INTRODUCTION

Dysmenorrhea is defined as cramp-like pain in the lower abdomen during menstruation without underlying pathology (primary dysmenorrhea) and is a highly prevalent gynecological condition among adolescent females. Pain typically begins shortly before or at the onset of menses and can last several days, affecting school attendance, participation in activities, and psychological well-being. Epidemiological studies estimate prevalence rates between 50% and 90% among adolescents worldwide, with up to 10–20% experiencing severe symptoms that disrupt daily life.

Conventional therapeutic approaches primarily rely on NSAIDs such as ibuprofen or mefenamic acid, which reduce prostaglandin synthesis to alleviate cramping. However, gastrointestinal side effects, contraindications, and individual preferences for natural remedies have driven research into alternative and complementary therapies. Herbal extracts with analgesic activity are increasingly studied, among which pudhina (mint) is promising due to its wide availability, traditional use, and favorable safety profile.

2. *Pudhina* (Mint): Botanical Profile and Bioactive Compounds

Pudhina commonly referred to as peppermint (*Mentha piperita*) or related *Mentha* species is a widely cultivated aromatic herb used traditionally in culinary, medicinal, and therapeutic contexts. The plant's pharmacological effects are attributed mainly to its essential oil components, particularly **menthol**, along with menthone and various flavonoids. Menthol interacts with transient receptor potential (TRP) cation channels such as TRPM8, producing a cooling sensation that is linked to analgesic effects and modulation of nerve conduction related to pain perception. In addition, mint exhibits antispasmodic effects on smooth muscles and anti-inflammatory activity, which are relevant to uterine cramping and associated menstrual discomfort.

Mechanistically, these compounds may inhibit prostaglandin synthesis similar to NSAIDs reduce calcium influx in smooth muscle, and influence endogenous opioid pathways, contributing to decreased pain perception. The aromatic compounds may also exert central analgesic effects and reduce associated symptoms such as nausea and headaches. These multimodal actions make pudhina extract a candidate for menstrual pain management, particularly in natural or holistic health care frameworks.

3. Evidence Review: Clinical and Experimental Studies

3.1 Randomized Controlled Trials

A key clinical investigation compared the efficacy of peppermint capsules with mefenamic acid a standard NSAID in young women with primary dysmenorrhea. In this double-blind randomized crossover trial, peppermint extract significantly reduced pain intensity and duration of cramps, although mefenamic acid showed slightly greater reduction in bleeding volume and overall analgesic use. Participants in the mint group reported fewer gastrointestinal side effects such as nausea and diarrhea, highlighting a potential tolerability advantage of herbal intervention.

Another randomized double-blind clinical trial evaluating 330 mg peppermint capsules taken three times daily during menstruation found a statistically significant reduction in pain severity compared to placebo in young women with primary dysmenorrhea. The study suggests that menthol-rich capsules could be effective in moderating menstrual pain severity, although pain duration did not differ significantly between groups.

3.2 Systematic Evidence

A systematic review of randomized controlled trials on peppermint's effects on menstrual disorders, including dysmenorrhea and related symptoms, found that peppermint may lead to statistically significant reductions in pain severity and improvements in associated cognitive and gastrointestinal symptoms. However, the authors emphasized limitations in study quality and consistency, calling for more rigorous trials to confirm effectiveness and clarify mechanisms.

3.3 Adolescent-Focused Studies

Specific research targeting adolescents is relatively limited but emerging. A quasi-experimental study among teenage girls found that mint leaf extract significantly reduced menstrual pain levels compared with a control group without treatment, indicating positive outcomes in adolescent populations. Another descriptive intervention study reported lowered dysmenorrhea symptoms following daily mint extract consumption, although methodological limitations such as non-randomized design and small sample sizes were noted.

4. Potential Mechanisms of Action in Dysmenorrhea

Mint's analgesic and antispasmodic effects are largely attributed to menthol's interaction with TRPM8 receptors, which produce sensory "cooling" and may modulate nociceptive pathways involved in menstrual pain. This interaction triggers calcium release from intracellular stores and influences neuronal activity related to pain perception. Additionally, menthol and other bioactive compounds in mint may inhibit the production of pro-inflammatory cytokines, reduce smooth muscle contraction through calcium channel modulation, and activate endogenous opioid pathways, collectively contributing to pain relief and muscle relaxation.

These pharmacological pathways align with the pathophysiology of primary dysmenorrhea, which is mediated by prostaglandin-induced uterine hypercontractility and inflammation. By targeting multiple aspects of this pathophysiology, pudhina extract offers a multimodal approach to symptom reduction.

5. Practical Implications for Adolescent Health

5.1 Efficacy and Safety

Based on current clinical evidence, pudhina extract appears to reduce the intensity of menstrual pain and associated symptoms when used during menstruation. The favorable side effect profile, ease of administration (e.g., capsules, teas, or extracts), and cultural acceptance of mint as a traditional remedy may support its use as a complementary therapy, especially in adolescents who may be sensitive to NSAID-related side effects.

5.2 Dosage and Administration

While clinical trials have used defined dosages (e.g., 330 mg peppermint capsules three times daily), standardized dosing guidelines remain undeveloped. Variability in herb quality, formulation, and peppermint chemotype can affect bioavailability and therapeutic outcomes. Health practitioners should consider these factors and exercise caution, especially in adolescent populations, until consensus dosing and regulatory standards are established.

5.3 Integration into Holistic Care

Incorporating pudhina extract into a broader, evidence-based dysmenorrhea management plan—including lifestyle modifications, dietary adjustments, stress management, and physical therapies—can support comprehensive care. Education about realistic expectations, possible interactions with other medications, and awareness of individual responses is essential when recommending herbal interventions.

6. Limitations and Future Research

Despite encouraging findings, limitations remain. Many studies involve small sample sizes, mixed age groups, and non-standardized formulations. Research in adolescent-specific cohorts is particularly limited. Future investigations should prioritize:

- **Large-scale randomized controlled trials** focusing on adolescents with standardized pudhina extract.
- **Dose-response studies** to identify optimal therapeutic ranges for pain relief.
- **Mechanistic research** using biochemical markers to elucidate pathways of action.
- **Comparative studies** with existing first-line therapies to establish relative efficacy and safety.

Such studies will enhance clinical confidence in recommending pudhina extract and clarify its role in evidence-based dysmenorrhea management.

7. Conclusion

Pudhina (*Mentha* spp.) extract, rich in menthol and other bioactive compounds, shows promise in alleviating dysmenorrhea symptoms among adolescent girls and young women. Clinical studies indicate reductions in pain severity and improvements in related symptoms, with a favorable safety profile compared to conventional NSAIDs. However, limitations in study design and heterogeneity in formulations highlight the need for further robust research. Integrating pudhina extract into holistic menstrual health strategies may offer a valuable, low-risk option for adolescents seeking alternative or complementary relief from menstrual pain.

8. References

1. Fitriah, M. S., Kusnianto, H., Sari, Y. K., & Handayani, H. (Year). *Effect of peppermint therapy on the level of primary dysmenorrhea pain: A systematic literature review. Jurnal Ners dan Kebidanan*.
2. Heshmati, A., Dolatian, M., Mojab, F., & Nikkhah, S. (2016). *The effect of peppermint (Mentha piperita) capsules on the severity of primary dysmenorrhea*. *Journal of Herbal Medicine, 6*(3), 137–141. https://doi.org/10.1016/j.hermed.2016.05.001
3. Saraf, P., Davasir Vadam, V., & Kaur, R. (2022). *Effectiveness of mint extract on dysmenorrhea related to pre-menstrual symptoms among adolescent girls*. *International Journal of Applied Research, 8*(6), 589–591.
4. Systematic Review on Effects of Peppermint on Menstrual Disorders. (2025). *Iranian Journal of Nursing and Midwifery Research*.
5. Wijayanti, E., Ernawati, & Arista Apriani. (2020). *Effectiveness of mint leaf extract (Mentha piperita Linn.) on menstrual pain level in adolescents*. *Advances in Health Sciences Research*, 1st International Conference on Community Health.