



EFFECTS OF TOBACCO ON PERIODONTIUM –AN UPDATE

Aarti Charokar, PhD Scholar, Department of Microbiology, RKDF Dental College & Research Centre, Bhopal.

Dr.AbhishakGupta, PhD Guide. SRK University, Bhopal

Abstract

Tobacco is used as a psychoactive substance. Tobacco smoke anticipated to incorporate over 4000 compounds lots of which can be pharmacologically active, toxic, mutagen and carcinogenic. Nicotine is one of the most studied additives of tobacco products and the maximum pharmacologically lively compound in tobacco smoke. Nicotine is a poisonous alkaloid determined in tobacco smoke and can input the body via absorption thru the oral mucosa and skin or inhalation via the lungs. Nicotine is likewise relatively addictive. Only 2.5 percent of the people who smoke try and end smoking are successful.

Periodontal disease afflicts many people in some unspecified time in the future during existence and is a chief motive of tooth loss in adults. The distributions of Periodontitis (inflammation of the periodontium) within the population suggest that a subset of individuals is exceedingly at risk of this infection, even as closing majority exhibits various ranges of resistance and slight susceptibility. There's evidence that cigarette smoking exerts both systemic and nearby results, though research had proven an affiliation between the 2, specific mechanisms underlying the consequences of smoking on periodontal reputation and wound recovery continue to be unresolved.

Keywords: Periodontal disease, Tobacco, Lungs, nicotine, mucosa

Introduction

Inflammation confined only to the marginal gingiva is known as gingivitis. Once it extends to involve the connective tissue attachment and alveolar bone loss, it is called periodontitis. Chronic periodontitis causes more teeth loss than any other disease for adult's does. Periodontal disease afflicts many individuals at some point during life and is a major cause of tooth loss in adults. The distribution of Periodontitis (inflammation of the periodontium) in the population suggest that a subset of individuals is highly susceptible to this infection, while remaining majority exhibit varying degrees of resistance and moderate susceptibility. Current data indicate that dental plaque is the cause of Periodontitis and cigarette smoking is a known risk factor, which adversely influences periodontal health. There is evidence that cigarette

smoking exerts both systemic and local effects. Though studies had shown an association between the two, precise mechanisms underlying the effects of smoking on periodontal status and wound healing remain unresolved. The disease is initially confined to the gingival and is called chronic marginal gingivitis; later the supporting tissues are involved, and are called marginal periodontitis. Gingivitis is a problem frequently occurring in young adults, which often gets unnoticed.¹

Periodontitis or inflammation of the periodontium result from the extension of the inflammatory process initiates the gingival to the supporting periodontal tissues in which case there is bone damage and loosening and eventual loss of teeth. The Periodontium is a connective tissue organ, covered by epithelium that attaches the teeth to the bones of the jaws and provides a continually adapting apparatus for support of teeth during function. The periodontium comprises of four connective tissues. They are the cementum, alveolar bone, periodontal ligament and the gingiva. The gingiva is a part of the oral mucosa that covers the alveolar process of the jaws and surrounds the neck of the tooth. The periodontal ligament is the connective tissue structure that surrounds the root and connects it to the bone. The cementum is a calcified tissue that forms the outer covering of the anatomical root of the teeth. The alveolar process is the bone that forms and supports the tooth sockets.

Constituents of Tobacco

Tobacco is used as a psychoactive substance. Tobacco smoke estimated to contain over 4000 compounds many of which are pharmacologically active, toxic, mutagen and carcinogenic. There are 43 known carcinogens in tobacco smoke.³ Tobacco smoke consists of a gaseous phase which is composed of carbon monoxide, nitrogen, oxygen, carbon dioxide; tar as well as particulate phase contains nicotine, water and polycyclic aromatic hydrocarbons.⁴

Nicotine

Nicotine, is the second most abundant constituent of tobacco smoke, is responsible for the addictive properties of tobacco. It's an addictive substance which release dopamine—a chemical in the brain that is associated with the feeling of pleasure.⁵ Nicotine is one of the most studied components of tobacco products and the most pharmacologically active compound in tobacco smoke. It has been shown to have various mood altering effects on its consumers. Nicotine is a poisonous alkaloid found in tobacco smoke and can enter the body by absorption through the oral mucosa and skin or inhalation via the lungs.⁶

Carbonmonoxide

The amount of oxygen carried by the blood may be severely deprived in heavy smokers due to the effects of carbon monoxide. Oxygen levels may be reduced by as much as 15%. Oxygen level reduced due to carbon monoxide which effect fetus, low weight of babies born to women who smoke.³

Nitrogenoxide

Nitrous oxide gases are formed by the combustion of nitrogen- containing amino acids and proteins in the tobacco leaf. Damage the respiratory airways. When smokers release nitrogenoxide in air they cause many effect to environment. Breathing high level of nitrogen oxide can cause swelling of tissue in the throat and upper respiratory tract, reduced oxygenation of body tissue.

Hydrogencyanide

The volatile nitriles, such as HCN, acetonitrile, and acrylonitrile, are important because of their toxic effects. Hydrogen cyanide is poisonous gases which are formed from the combustion of the protein and nitrate compounds existed in tobacco. They effect neurological, respiratory, cardiovascular and thyroid. They are not carcinogen but affect other health problem. HCN is a poisonous gas which leads to problem of central nervous system.

Radioactive compounds

The radio active compounds found in highest concentration in cigarette smoke are polonium-210 and potassium-40. Radioactive compounds are well established as carcinogens.³ Metals – Thirty metals have been detected in tobacco smoke including nickel, arsenic, cadmium, chromium and lead.³

Oral and systemic effects of tobacco

Tobacco is the most important etiological factor for oral diseases including oral cancer, oral mucosal lesions and periodontal disease. Tobacco affects both local and systemic either it is used in the form of chewing and smoking.

Effects of smokeless tobacco (chewers)

Tobacco was most often chewed as an ingredient in betel quid (smokeless tobacco or paan) in India. Chewers or person who takes tobacco in the form of snuff and ghutka. Oral effect of Smokeless tobacco is typically seen on the mucosal surface where the product is placed, as well as periodontium. The affected site or lesion can be white or yellowish-brown in color and it is known as leukoplakia. Smokeless tobacco contains high levels of sodium which may contribute to elevated blood pressure and heart rate. The most consistently reported periodontal effects of Smokeless tobacco (Chewers) are mucosal lesions that may also affect the gingiva, can cause gingival recession and various form of periodontal diseases.

Smokeless tobacco which provides nicotine along with other substance that is known to cause cancer. While it is not inhaling directly to lungs, it is still being absorbed through the lining of mouth and circulating through blood stream to all parts of the body, and cause bacteremia and endotoxemia, which is responsible for damage of the vascular endothelial integrity, platelet function and blood coagulation. Use of smokeless tobacco has linked to oral cancer, esophageal cancer, and pancreatic cancer. It can also cause heart disease, oral lesion (leukoplakia) and gum disease (periodontitis).

Effect of smokers

For many years, smoking has been linked with lung disease, cancer, cardiovascular disease and poor pregnancy outcomes, such as miscarriage and low birthweight.⁷ Over the past decades, it has also been recognized that smoking is associated with periodontal disease. As early as the 1940s, Pindborg noted that acute necrotizing ulcerative was associated with smoking in the Danish Royal Marines.⁸

Tobacco as a risk factor for periodontal disease

(Effect of tobacco with periodontal disease)

An individual's periodontal health has been identified to be affected by tobacco chewing and smoking. Tobacco intake is related with increasing rate of multiple health complications which also includes loss of periodontal bone and the loss of the periodontal attachment and bone formation. In addition, tobacco smoking and chewing has been connected with the masking of the gingival swelling symptoms: Gingivitis is a disease that has been related with smoking for a long time.⁹ Studies have revealed that increased rate of tobacco smoking and gingival bleeding on probing are the early symptoms of gingivitis, a subject that is not dependent on significant signs of gingivitis and is now broadly applied towards identifying lesions in periodontal complications.¹⁰ Gingival bleeding is higher in non-smokers than in smokers.¹¹ This is attributed to the fact that there is vasoconstriction of the gingival cells or there is an occurrence of the keratinisation of the gingival among the smokers.¹²

Periodontal diseases are a group of conditions affecting the supporting structures of the dentition. Progression and severity of the disease depends on the complexity reactions between the host defence mechanism and risk factor such as microbial growth, genetic and environmental factor and oral hygiene as well as age, sex and race. Tobacco smoking is a significant risk factor for periodontal disease.¹³ Tobacco smoking is an addictive habit first introduced into Europe. Smokers are three times more likely to get a cute Periodontitis than non-smokers.¹⁴

Periodontitis start from gingivitis (gum disease) where specific bacteria present in plaque play important role to cause gingivitis. Then gingivitis or bacteria resulting in progressive destruction of periodontal ligament and alveolar bone with pocket formation, and in periodontal pocket anaerobic bacteria grow and enter in blood stream and cause bacteremia and endotoxemia.

Prevalence of plaque development in smokers

Cigarette smoking alone does not cause Periodontitis disease. Periodontal disease is caused by bacterial Plaque that gets under the gingival tissue and initiates bone loss around the teeth. Smokers show a higher prevalence of dental plaque than non-smokers suggested that more severe periodontal disease in smokers might be because of greater accumulation of plaque which play important role to cause periodontal disease.

Prevalence of probing depth in smokers

On the premise of CPITN: The difference between people who smoke and non-smokers in probing intensity can also be an evidence as smokers have deeper periodontal pockets than non-people who smoke, which might confound the effect of smoking on periodontal pathogens (Kigure et al 1995) Stoltenberg et al²⁴ reported that the chances ratio for having a mean probing intensity ≥ 3.5 mm was five three times more in people who smoke. Stoltenberg et al²⁴ located that smoking changed into the most powerful risk indicator for extended pocket intensity.

Early onset of inflammation in smokers

Smokers also had an earlier onset of clinically visible inflammation compared to non-smokers, which was attributed to the early pathogenic colonization, leading to sustained pathogen enrichment with periodontal pathogens. This is consistent with other findings in smokers, indicating that gingivitis is preceded by a decrease in the abundance of early colonizers, such as the genera *Streptococcus* and *Veillonella*, and an increase in the abundance of periodonto-pathogens, such as the Genera *Treponema* and *Selenomonas* (Matthewsetal.,2013;Peruzzoet al., 2016).¹⁵

Smoking and microorganism

Haffajee and Socransky¹⁶ investigated the connection between cigarette smoking and subgingival microbiota the usage of checkerboard DNA hybridization. Kamma and colleagues¹⁷ investigated the consequences of smoking in topics with aggressive periodontitis. They said that smokers harbored more quantity of strict anaerobes, specially *Porphyromonas gingivalis*, *Bacteroides forsythus* (*Tanarellaforsythus*) and *Eikenella corrodens* compared to non-smoking topics.²

Tobacco smoke carries phenols and cyanides, that may account for antibacterial and toxicproperties.Smoker harboured extensively higher levels and have been at significantly more threat of infection with *Tanarella forsythus* than non-people who smoke.¹⁸ Adjusting for illnesses verity, *Porphyromonas gingivalis* turned into also more likely to subgingivally infect people who smoke than non-smokers.¹⁹ opinions have been divided about the effect of smoking on continual inflammatory periodontal disease. earlier reviews of the epidemiology of periodontal disorder concluded that smoking was a possible causative factor. Few research have conclusively demonstrated any applicable microbiological adjustments inside the periodontal tissues due to smoking. a few authors the use of self-pronounced smoking information investigated the relationship between periodontal pathogens and cigarette intake. They reported an increased hazard for smokers to have subgingival infection with *Porphyromonas gingivalis* even though this became no longer found to be statistically sizeable. in this identical have a look at the investigators found people who smoke were 3times much more likely to harbor a *actinomycetemcomitans*.^{20,21} Many authors investigated the connection among cigarette smoking and the superiority of periodontal pathogens the use of polymerase chain reaction strategies. on this examine, which covered same numbers of smoking and non-smoking topics with generalized competitive periodontitis, the investigators should discover no good sized variations in the occurrence of any of the pathogenic species which blanketed *Porphyromonas gingivalis*, *Prevotella intermedia*, *Tanarella forsythus*, *Actinobacillus actinomycetemcomitans* and *T.denticola*.^{22,23}

References

1. QuantidIR, Sandhu HS, Mathews DC. "Tobacco smoking and periodontal disease". Journal of Canadian Dental Association 1997, March; 63(3): 187-92,194-5.
2. Tarulatha R ShyPreagali, Mohamed Helmy Salama, Deepak P Bhayya, Prevalence of tobacco usage and its effect on the periodontal health parameters in the mining employees and the general population- A comparative study, BJOS, volume 16, 2017
3. Sobnen Peter Fourth Edition Essentials of community and Preventive dentistry Arya Publication House 2009. 135-4.
4. McGuire JR et al cotinine in Saliva and Gingival Crevicular fluid of Smokers with Periodontal Disease. J Periodontal 1989; 176-81. <http://dx.doi.org/10.1902/jop.1989.60.4.176> PMID2656979
5. Benowitz NL, Jacob P, Jones RT, Rosenberg J, " Interindividual variability in the metabolism and Cardiovascular effects of nicotine in man" J pharmacol Exp Ther 1982,221 368-72 Pmid.
6. Holt SC, Ebersole JL. Porphyromonas gingivalis, Treponema denticola, and Tannerella forsythia: The " red complex ", a prototype polybacterial pathogenic consortium in periodontitis. Periodontology 2000. 2005; 38: 72-122.
7. Wald, N.J. and Hackshaw , A.K. (1996) Cigarette smoking : An epidemiological overview, British Medical Bulletin , 52, 3-11.
8. Pindborg, J.J. (1949) Tobacco and gingivitis ; correlation between consumption of tobacco, ulceromembranous gingivitis and calculus. Journal of Dental Research,28 460-463.<http://dx.doi.org/10.1177/00220345490280050601>
9. Herman Al.Sofuoglu M Cognitive effects of nicotine genetic moderators Addict Biol 2010 15 (3) 250-65 DOI 10.1111/j 1369-1600 2010 002132
10. Hirschfelder AB Tobacco Santa Barbara Calif Greenwood 2010 I -264.
11. Marijia Ivic-Kardum, Natasa Belder et al Diagnostic Methods of Evaluation of Microbial flora in Periodontitis, Acta Stomatol Croat, Vol. 35, br. 1, 2001
12. Kuramitsu, H.K. (1998). Proteases of Porphyromonas gingivalis: what don't they do? Oral Microbiology and Immunology, Vol.13 (No.5):263-27
13. Ojima M, Hanioka T Destructive effect of smoking on molecular & genetic factor of periodontal disease. Tobacco induced disease 2010; 8: 4.

14. SS Hiremath Textbook of preventive and community Dentistry Elsevier 2007. 129
15. Clarke JK. On the bacterial factor in the etiology of dental caries. *British Journal of Experimental Pathology*. 1924;5: 141-147
16. Mitchell DF, Johnson M. The nature of the gingival plaque in the hamster: Production, prevention and removal. *Journal of Dental Research*. 1956; 35:651-655.
17. Socransky S, Macdonald JB, Sawyer S. The cultivation of *Treponema microdentium* as surface colonies. *Archives of Oral Biology*. 1959; 1: 171-172
18. Zambon JJ, Grossi SG, Machtei EE, Ho A, Dunford R, Genco J. Cigarette smoking increases the risk for subgingival infection with periodontal pathogens. *J Perio* 1996; 67: 1050- 1054.
19. Sayers NM, James JA, Drucker DB, Blinkhorn AS. Possible potentiation of toxins from *Prevotella intermedia*, *Prevotella nigrescens* and *Porphyromonas gingivalis* by cotinine *J perio* 1999; 70: 1269.
20. Mager DL, Haffajee AD, Socransky SS. Effects of periodontitis and smoking on the microbiota of oral mucous membranes and saliva in systemically healthy subjects *J Clin Perio* 2003;30:1031-1037
21. Luciana Machion L, Andia DC, Saito D et al . Microbiological changes with the use of Locally Delivered Doxycycline in the periodontal Treatment of Smokers. *J Perio* 2004; 75:1600-1604.
22. Bostrom, L., Bergstrom, J., Dahlen, G., and Linder, L. E. (2001). Smoking and subgingival microflora in periodontal disease. *J. Clin. Periodontol.* 28, 212–219.
23. Kamma JJ, Nakou M, Baehni PC. Clinical and microbiological characteristics of smokers with early onset periodontitis. *J Perio Res* 1999; 34: 25 – 33.
24. Stoltenberg JL, Osborn JB, Pihlstrom BL, Herzberg MC, Aepli DM, Wolff LF, Fischer GE: Association between cigarette smoking, bacterial pathogens, and periodontal status. *J Periodontol* 1993, 64(12):1225-123